

# Science Summary

## Dairy and Inflammation



### Overview

Dairy foods such as milk, cheese and yogurt are foundational foods in healthy dietary patterns recommended by the Dietary Guidelines for Americans (DGA). Healthy dietary patterns that include dairy foods are linked with lower risk of key chronic diseases such as type 2 diabetes, cardiovascular disease and obesity. These noncommunicable diseases are sometimes called “inflammatory diseases,” because they often co-occur with chronic, systemic inflammation. Lifestyle factors such as diet may impact chronic inflammation. Emerging evidence indicates that consuming dairy foods, including whole- and reduced-fat dairy foods, is not linked to increased levels of inflammatory markers. Some research indicates that consuming certain dairy foods may be linked to lower levels of some inflammatory markers.

### Healthy dietary patterns with dairy can help lower risk for chronic diseases linked with inflammation

Chronic diseases like cardiovascular disease (CVD), type 2 diabetes (T2D) and obesity affect millions of Americans and result in high healthcare costs and lost productivity.<sup>1-3</sup> These conditions are also associated with higher levels of inflammatory markers.<sup>4,5</sup> A healthy diet is the foundation for prevention and management of several chronic diseases, including CVD, T2D and obesity.<sup>5-10</sup> The Scientific Report of the 2020 Dietary Guidelines Advisory Committee emphasizes this point, stating that risk factors for CVD, such as inflammatory markers, are “favorably influenced by habitual adherence to dietary patterns that include fruits, vegetables, whole grains, legumes, nuts, unsaturated vegetable oils, fish, seafood, [and] lower fat dairy products.”<sup>11</sup> The Healthy U.S.-Style Dietary Pattern in the 2020 DGA includes these foods, recommending 3 daily servings of low-fat or fat-free dairy foods for those 9 years and older, 2½ servings for children 4-8 years and 2 servings for children 2-3 years.<sup>10</sup> It also recommends 1⅓ to 2 servings of whole- and reduced-fat dairy foods for toddlers 12-23 months and small amounts of yogurt and cheese for infants 6 to 12 months, depending on developmental readiness.<sup>10</sup>

### Eating dairy foods is not linked to inflammatory markers in systematic reviews or meta-analyses

Eating dairy foods does not increase levels of inflammatory biomarkers in individuals without a milk protein allergy, according to results of two meta-analyses and three systematic reviews. Six of the randomized controlled trials of healthy adults included in a meta-analysis by Benatar et al.<sup>12-18</sup> assessed the impact of low-fat or high-fat dairy foods on levels of C-reactive protein (CRP), a commonly used biomarker of inflammation produced by the liver. There were no differences in CRP levels between the dairy-rich diets and the control diets in four of these studies, even when higher-fat dairy foods were included in the dairy intervention. Zemel et al. reported lower levels of CRP at the end of a 28-day study,<sup>16</sup> and Stancliffe et al. also reported significantly lower CRP levels in the group consuming 3.5 daily servings of dairy foods.<sup>14</sup> Another meta-analysis used an “inflammatory score,” which combines many biomarkers

of inflammation into a single metric, to evaluate the results of 52 trials on dairy intake and inflammation.<sup>19</sup> This meta-analysis concluded that diets containing dairy foods, especially fermented dairy foods, had a modest anti-inflammatory effect most apparent in individuals with metabolic disorders like obesity. Three systematic reviews reported similar results, concluding that eating dairy foods had a beneficial impact or no impact on levels of inflammatory markers in both healthy individuals<sup>20,21</sup> and those with metabolic disorders.<sup>20-22</sup>

---

### **Clinical trials find that consuming dairy foods exerts no change or favorable change on some biomarkers of inflammation**

In addition to the studies covered in meta-analyses and systematic reviews, at least five additional randomized controlled trials<sup>23-27</sup> have been published on the links between dairy foods and inflammation. These studies provide additional evidence that eating dairy foods does not cause inflammation in adults.

In one study, 139 adults with risk factors for metabolic syndrome ate either ~3 ounces of whole-fat cheese, low-fat cheese or a bread and jam control with the same amount of calories for 12 weeks.<sup>23</sup> There were no differences in CRP levels between these groups, indicating that a dairy-rich diet, even with whole-fat cheese, was not linked with increased inflammation. In another study<sup>24</sup> completed by 52 middle-aged adults, participants consumed an energy-deficit diet providing either high or low amounts of dairy foods for 24 weeks. There were no differences in CRP levels between the two groups after the intervention. In a similar randomized crossover study,<sup>25</sup> 45 middle-aged, overweight men and postmenopausal women consumed a high-dairy diet (5-6 daily servings of dairy foods) or a low-dairy diet (<1 daily serving of dairy foods) for 6 weeks. There was no significant difference in CRP levels between the two groups following the intervention. A randomized crossover trial of 92 men and women with abdominal obesity also found no differences in CRP levels after participants ate a diet rich in saturated fat from cheese, a diet rich in saturated fat from butter, a diet rich in monounsaturated fatty acids, a diet rich in polyunsaturated fatty acids, or a low-fat high carbohydrate diet for 4 weeks.<sup>27</sup> Another randomized controlled trial assessed the impact of eating yogurt on markers of chronic inflammation in healthy women with and without obesity.<sup>26</sup> Women consumed 12 ounces of either low-fat yogurt or soy pudding for 9 weeks. Eating yogurt led to decreased levels of several, but not all, markers of inflammation measured in this study in both the women with obesity and the women without obesity.

---

### **Dairy foods are scored as neutral or “anti-inflammatory” by dietary inflammation scoring system**

A new, validated approach to assessing dietary and lifestyle factors that affect systemic inflammation was published in 2019.<sup>28</sup> The research team used food frequency questionnaires and levels of inflammatory biomarkers to assess dietary intake and create an inflammation biomarker score. They validated this score with 3 study populations including over 14,000 Americans. Both high-fat dairy foods (including whole- and reduced-fat milk, yogurt, cheese and ice cream) and low-fat dairy foods (including low- and fat-free milk, low-fat cheeses and ice cream) received negative scores (-0.14 and -0.12, respectively), indicating an anti-inflammatory effect. This scoring system, in conjunction with the other published literature on the topic, indicates that dairy foods do not cause inflammation and are either neutral or mildly anti-inflammatory.

## References

- <sup>1</sup> Adult Obesity Facts | Overweight & Obesity | CDC. <https://www.cdc.gov/obesity/data/adult.html>. Accessed November 20, 2019.
- <sup>2</sup> Centers for Disease Control and Prevention. Division of Diabetes Translation At A Glance. <https://www.cdc.gov/chronicdisease/resources/publications/aag/diabetes.htm>. Published 2019.
- <sup>3</sup> Benjamin EJ, Muntner P, Alonso A, et al. Heart Disease and Stroke Statistics-2019 Update: A Report From the American Heart Association. *Circulation*. 2019;139(10):e56-e528. doi:10.1161/CIR.0000000000000659
- <sup>4</sup> Inflammation and Heart Disease | American Heart Association. <https://www.heart.org/en/health-topics/consumer-healthcare/what-is-cardiovascular-disease/inflammation-and-heart-disease>. Accessed November 17, 2020.
- <sup>5</sup> McLaughlin T, Ackerman SE, Shen L, Engleman E. Role of innate and adaptive immunity in obesity-associated metabolic disease. *J Clin Invest*. 2017;127(1):5-13. doi:10.1172/JCI88876
- <sup>6</sup> Diabetes Basics | CDC. <https://www.cdc.gov/diabetes/basics/>. Accessed August 19, 2020.
- <sup>7</sup> Eckel RH, Jakicic JM, Ard JD, et al. 2013 AHA/ACC Guideline on Lifestyle Management to Reduce Cardiovascular Risk: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2014;63(25 Pt B):2960-2984. doi:10.1016/j.jacc.2013.11.003
- <sup>8</sup> Van Horn L, Carson JAS, Appel LJ, et al. Recommended Dietary Pattern to Achieve Adherence to the American Heart Association/American College of Cardiology (AHA/ACC) Guidelines: A Scientific Statement From the American Heart Association. *Circulation*. 2016;134(22). doi:10.1161/CIR.0000000000000462
- <sup>9</sup> CDC. Adult Obesity Facts | Data | Adult | Obesity | DNPAO | CDC. <http://www.cdc.gov/obesity/data/adult.html>. Published 2016. Accessed December 4, 2015.
- <sup>10</sup> USDA and HHS. 2020-2025 Dietary Guidelines for Americans.; 2020. [https://www.dietaryguidelines.gov/sites/default/files/2020-12/Dietary\\_Guidelines\\_for\\_Americans\\_2020-2025.pdf](https://www.dietaryguidelines.gov/sites/default/files/2020-12/Dietary_Guidelines_for_Americans_2020-2025.pdf).
- <sup>11</sup> Dietary Guidelines Advisory Committee. 2020. Scientific Report of the 2020 Dietary Guidelines Advisory Committee. [https://www.dietaryguidelines.gov/sites/default/files/2020-07/ScientificReport\\_of\\_the\\_2020DietaryGuidelinesAdvisoryCommittee\\_first-print.pdf](https://www.dietaryguidelines.gov/sites/default/files/2020-07/ScientificReport_of_the_2020DietaryGuidelinesAdvisoryCommittee_first-print.pdf).
- <sup>12</sup> Benatar JR, Sidhu K, Stewart RAH. Effects of High and Low Fat Dairy Food on Cardio-Metabolic Risk Factors: A Meta-Analysis of Randomized Studies. *PLoS One*. 2013;8(10):1-12. doi:10.1371/journal.pone.0076480
- <sup>13</sup> G.E. C, P.R.C. H, J.D. B, A.M. C, K.J. M. Dairy consumption and cardiometabolic health: Outcomes of a 12-month crossover trial. *Nutr Metab*. 2012;9:1-11. doi:10.1186/1743-7075-9-19
- <sup>14</sup> Stancliffe RA, Thorpe T, Zemel MB. Dairy attenuates oxidative and inflammatory stress in metabolic syndrome. *Am J Clin Nutr*. 2011;94(2):422-430. doi:10.3945/ajcn.111.013342
- <sup>15</sup> Wennersberg MH, Smedman A, Turpeinen AM, et al. Dairy products and metabolic effects in overweight men and women: results from a 6-month intervention study. *Am J Clin Nutr*. 2009;90(4):960-968. doi:10.3945/ajcn.2009.27664
- <sup>16</sup> Zemel MB, Sun X, Sobhani T, Wilson B. Effects of dairy compared with soy on oxidative and inflammatory stress in overweight and obese subjects. *Am J Clin Nutr*. 2010;91(1):16-22. doi:10.3945/ajcn.2009.28468
- <sup>17</sup> van Meijl LEC, Mensink RP. Effects of low-fat dairy consumption on markers of low-grade systemic inflammation and endothelial function in overweight and obese subjects: an intervention study. *Br J Nutr*. 2010;104(10):1523-1527. doi:10.1017/S0007114510002515
- <sup>18</sup> Benatar JR, Jones E, White H, Stewart RAH. A randomized trial evaluating the effects of change in dairy food consumption on cardio-metabolic risk factors. *Eur J Prev Cardiol*. 2014;21(11):1376-1386. doi:10.1177/2047487313493567
- <sup>19</sup> Bordoni A, Danesi F, Dardevet D, et al. Dairy products and inflammation: A review of the clinical evidence. *Crit Rev Food Sci Nutr*. 2017;57(12):2497-2525. doi:10.1080/10408398.2014.967385
- <sup>20</sup> Ulven SM, Holven KB, Gil A, Rangel-Huerta OD. Milk and Dairy Product Consumption and Inflammatory Biomarkers: An Updated Systematic Review of Randomized Clinical Trials. *Adv Nutr*. 2019;10(suppl\_2):S239-S250. doi:10.1093/advances/nmy072
- <sup>21</sup> Nieman KM, Anderson BD, Cifelli CJ. The Effects of Dairy Product and Dairy Protein Intake on Inflammation: A Systematic Review of the Literature. 2020. doi:10.1080/07315724.2020.1800532
- <sup>22</sup> Labonté M-È, Couture P, Richard C, Desroches S, Lamarche B. Impact of dairy products on biomarkers of inflammation: a systematic review of randomized controlled nutritional intervention studies in overweight and obese adults. *Am J Clin Nutr*. 2013;97(4):706-717. doi:10.3945/ajcn.112.052217
- <sup>23</sup> Raziani F, Tholstrup T, Kristensen MD, et al. High intake of regular-fat cheese compared with reduced-fat cheese does not affect LDL cholesterol or risk markers of the metabolic syndrome: a randomized controlled trial. *Am J Clin Nutr*. 2016;104(4):973-981. doi:10.3945/ajcn.116.134932
- <sup>24</sup> Bendtsen LQ, Blædel T, Holm JB, et al. High intake of dairy during energy restriction does not affect energy balance or the intestinal microflora compared with low dairy intake in overweight individuals in a randomized controlled trial. *Appl Physiol Nutr Metab*. 2018;43(1):1-10. doi:10.1139/apnm-2017-0234
- <sup>25</sup> Eelderink C, Rietsema S, Van Vliet IMY, et al. The effect of high compared with low dairy consumption on glucose metabolism, insulin sensitivity, and metabolic flexibility in overweight adults: A randomized crossover trial. *Am J Clin Nutr*. 2019;109(6):1555-1568. doi:10.1093/ajcn/nqz017
- <sup>26</sup> Pei R, DiMarco DM, Putt KK, et al. Low-fat yogurt consumption reduces biomarkers of chronic inflammation and inhibits markers of endotoxin exposure in healthy premenopausal women: a randomised controlled trial. *Br J Nutr*. November 2017;1-9. doi:10.1017/S0007114517003038
- <sup>27</sup> Brassard D, Tessier-Grenier M, Allaire J, et al. Comparison of the impact of SFAs from cheese and butter on cardiometabolic risk factors: A randomized controlled trial. *Am J Clin Nutr*. 2017;105(4):800-809. doi:10.3945/ajcn.116.150300
- <sup>28</sup> Byrd DA, Judd SE, Flanders D, Hartman TJ, Fedirko V, Bostick RM. The Journal of Nutrition Nutritional Epidemiology Development and Validation of Novel Dietary and Lifestyle Inflammation Scores. doi:10.1093/jn/nxz165