# Science Summary Dairy and Sarcopenia and Osteoporosis





## **Overview**

Including dairy foods in healthy dietary patterns can help people thrive at every age, including older adulthood. Eating dairy foods, a source of high-quality protein and other essential nutrients, may reduce the risk of age-related conditions such as sarcopenia (loss of muscle mass and strength) and frailty. Consuming dairy foods may also help older adults maintain healthy bones by reducing the risk of bone mineral density loss and lowering

the risk of osteoporosis-related falls and fractures. The 2020-2025 Dietary Guidelines for Americans (DGA) recommends that adults consume 3 servings of low-fat or fat-free dairy foods daily as part of the Healthy U.S.-Style Dietary Pattern.

# Dairy foods help support the nutritional needs of older adults

As the population of older adults in the U.S. continues to grow, the nutrition challenges these adults face will become even more serious public health concerns. Older adults may struggle with healthy eating due to agerelated changes and obstacles including socioeconomic factors, depression, changes in taste, physical ability to chew and swallow and body composition, along with reductions in appetite from medications and reduced mobility, among other challenges.<sup>1-5</sup> While men and women 70 years and older consume sufficient calories, on average, they do not consume enough of the nutrients of public health concern for Americans 2 years and older-calcium, vitamin D, fiber and potassium- and tend to exceed recommendations for daily sodium and saturated fat intake.<sup>6</sup> Older adults are also at risk for low intakes of protein and vitamin B12.<sup>6</sup>

Consuming dairy foods as part of healthy dietary patterns can help older adults thrive. Dairy foods like milk, yogurt and cheese are important sources of several shortfall nutrients for older adults, including protein, calcium, potassium, vitamin D and vitamin B12.<sup>7</sup> Dairy foods also provide high-quality nutrition at an affordable price<sup>8</sup> and are available in a variety of textures and flavors that may appeal to older adults experiencing physical challenges with eating. However, Americans 70 years and older consume, on average, less than 1½ servings of dairy foods per day.<sup>9</sup> The 2020 DGA recommends that all Americans 9 years and older consume 3 daily servings of low-fat or fat-free dairy foods in the Healthy U.S.-Style Dietary Pattern.<sup>6</sup> Adding at least one more daily serving of dairy foods to the dietary patterns of older adults can help bring them closer to meeting dairy recommendations.<sup>10,11</sup>



# Dairy foods are an important source of high-quality protein that can help meet the unique nutrition needs of older adults

Older adults have low intakes of protein relative to the Estimated Average Requirement (EAR),<sup>7</sup> and the 2020 DGA identifies dairy foods as a source of dietary protein for older adults.<sup>6</sup> Milk, cheese and yogurt are good sources of high-quality protein, which helps support bone health and preserve muscle mass and physical function in older adults as part of healthy dietary patterns.<sup>12</sup> Milk provides 8 grams (g) of high-quality protein per cup, and hard cheeses such as Cheddar and Colby provide 10 g or more of protein per 1½ ounce serving.<sup>13\*</sup> Some styles of yogurt like Greek yogurt can provide more than 20 g of protein per cup.<sup>13\*</sup> A modeling study examining the effect of increasing plant-based foods or dairy foods on protein intake in older adults found that doubling the intake of dairy foods could help older adults meet their protein needs. In contrast, doubling intake of plant-based foods resulted in an approximate 22 percent decline in protein consumption.<sup>14</sup> Because older adults currently do not meet dairy food recommendations, adding one more serving of dairy foods to their current diets would move them closer to both protein intake recommendations from the 2020 DGA.<sup>610,11</sup>

## Eating dairy foods may lessen risk of sarcopenia and frailty among older adults

Sarcopenia, or the loss of skeletal muscle strength and mass, accelerates with age, contributes to adverse health outcomes in older adults and may lead to substantial healthcare costs.<sup>15,16</sup> Frailty is also a common clinical syndrome in older adults, and it carries increased risks for poor health outcomes including institutionalization, falls, hospitalization and mortality.<sup>17</sup> Nutritional factors such as inadequate protein and energy intake play a role in the development of both sarcopenia and frailty.<sup>14,18-21</sup> Results of systematic reviews and prospective cohort studies indicate that consuming dairy foods like milk, cheese and yogurt in recommended amounts throughout older adulthood may help reduce the risk of sarcopenia and frailty by supporting muscle mass and muscle strength.

Results of two systematic reviews indicate that consuming dairy foods (low-fat milk, cheese and yogurt) may be beneficial for preserving muscle mass and muscle strength in adults 50 years and older, thereby reducing the risk of sarcopenia.<sup>15,22</sup> Two prospective cohort studies also reported similar results. One prospective cohort study found that consuming more than 7 servings per week of low-fat milk and yogurt versus less than 1 serving per week was linked with lower risk of frailty, slow walking speed and weight loss.<sup>23</sup> Another prospective cohort study concluded that consuming more meat and dairy foods was linked with a lower risk of developing physical frailty.<sup>24</sup>

#### Dairy foods may help support bone health in older adults

Eating dairy foods may also help older adults maintain healthy bones. Osteoporosis, the condition of porous, brittle and fragile bone, affects approximately 54 million Americans.<sup>25</sup> One out of every two postmenopausal women will experience an osteoporosis-related fracture at some point in her lifetime.<sup>26</sup> While some factors that affect peak bone mass are not modifiable (gender, ethnicity, genetics), others such as nutritional status, physical activity and body composition are modifiable. Unfortunately, older adults tend to fall short of meeting recommendations for nutrition needs associated with bone health like adequate calcium, vitamin D and potassium intake.<sup>7</sup>

\*FDC IDs: Cheddar: 1098009; Colby: 173416; Greek Yogurt: 1097564



Results of meta-analyses, a systematic review and longitudinal cohort studies indicate that eating dairy foods may be beneficial for bone health in older adults. Meta-analyses and systematic reviews as well as a prospective cohort study provide evidence that dairy foods can increase bone mineral density and reduce the risk of hip fracture.<sup>27-33</sup> A large systematic review with data from 91 publications concluded that moderate evidence indicates that daily intake of low-fat or fat-free dairy foods as part of a healthy dietary pattern may be associated with improved bone mineral density and fewer fractures in adults 50 years and older.<sup>32</sup> In two U.S. cohorts of approximately 43,000 men and 80,000 postmenopausal women aged 50 years and followed for up to 32 years, each 1 cup daily serving of milk was associated with an 8 percent lower risk of hip fracture.<sup>33</sup> Total dairy intake, about half of which was milk, was associated with a 6 percent lower risk of hip fracture per daily serving.<sup>33</sup> Some research reports that consuming dairy foods may not have an impact on certain indicators of bone health in older adults, such as bone mineral density, risk of falling and osteoporotic fractures.<sup>34-37</sup> These results indicate a need for continued research to further explain the links between dairy food intake and bone health in older adults.

#### References

- <sup>1</sup> Choosing Healthy Meals As You Get Older. https://www.nia.nih.gov/health/choosing-healthy-meals-you-get-older. Accessed December 3, 2019.
- <sup>2</sup> Walls AWG, Steele JG. The relationship between oral health and nutrition in older people. Mech Ageing Dev. 2004;125(12 SPEC.ISS.):853-857. doi:10.1016/j.mad.2004.07.011
- <sup>3</sup> Brownie S. Why are elderly individuals at risk of nutritional deficiency? Int J Nurs Pract. 2006;12(2):110-118. doi:10.1111/j.1440-172X.2006.00557.x
- <sup>4</sup> Morris H. Dysphagia in the elderly--a management challenge for nurses. Br J Nurs. 2006;15(10):558-562. doi:10.12968/bjon.2006.15.10.21132
- <sup>5</sup> Morley JE. Novel Approaches to Nutrition in Older Persons. Clin Geriatr Med. 2015;31(3):xiii-xiv. doi:10.1016/j.cger.2015.05.001
- <sup>6</sup> USDA and HHS. 2020-2025 Dietary Guidelines for Americans.; 2020. https://www.dietaryguidelines.gov/sites/default/files/2020-12/Dietary\_Guidelines\_ for\_Americans\_2020-2025.pdf.
- <sup>7</sup> Dietary Guidelines Advisory Committee. 2020. Scientific Report of the 2020 Dietary Guidelines Advisory Committee. https://www.dietaryguidelines.gov/ sites/default/files/2020-07/ScientificReport\_of\_the\_2020DietaryGuidelinesAdvisoryCommittee\_first-print.pdf.
- <sup>8</sup> Hess JM, Cifelli CJ, Agarwal S, Fulgoni VL. Comparing the cost of essential nutrients from different food sources in the American diet using NHANES 2011-2014. Nutr J. 2019;18(1):68. doi:10.1186/s12937-019-0496-5
- <sup>9</sup> National Dairy Council. NHANES 2015-2018. Hyattsville, MD; 2020.
- <sup>10</sup> Quann EE, Fulgoni VL, Auestad N. Consuming the daily recommended amounts of dairy products would reduce the prevalence of inadequate micronutrient intakes in the United States: diet modeling study based on NHANES 2007–2010. Nutr J. 2015;14(1):90. doi:10.1186/s12937-015-0057-5
- <sup>11</sup> Hess JM, Fulgoni VL, Radlowski EC. Modeling the Impact of Adding a Serving of Dairy Foods to the Healthy Mediterranean-Style Eating Pattern Recommended by the 2015-2020 Dietary Guidelines for Americans. J Am Coll Nutr. August 2018:1-9. doi:10.1080/07315724.2018.1485527
- <sup>12</sup> Phillips SM, Martinson W. Nutrient-rich, high-quality, protein-containing dairy foods in combination with exercise in aging persons to mitigate sarcopenia. Nutr Rev. 2019;77(4):2.16-2.29. doi:10.1093/nutrit/nuy062
- <sup>13</sup> USDA. FoodData Central. https://fdc.nal.usda.gov/index.html. Published 2019.
- <sup>14</sup> Houchins JA, Cifelli CJ, Demmer E, Fulgoni VL. Diet modeling in older Americans: The impact of increasing plant-based foods or dairy products on protein intake. J Nutr Health Aging. 2017;21(6):673-680. doi:10.1007/s12603-016-0819-6
- <sup>15</sup> Granic A, Dismore L, Hurst C, Robinson SM, Sayer AA. Myoprotective whole foods, muscle health and sarcopenia: A systematic review of observational and intervention studies in older adults. Nutrients. 2020;12(8):1-32. doi:10.3390/nu12082257
- <sup>16</sup> Bruyère O, Beaudart C, Ethgen O, Reginster JY, Locquet M. The health economics burden of sarcopenia: a systematic review. Maturitas. 2019;119:61-69. doi:10.1016/j.maturitas.2018.11.003
- <sup>17</sup> Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: Evidence for a phenotype. Journals Gerontol Ser A Biol Sci Med Sci. 2001;56(3). doi:10.1093/gerona/56.3.m146
- <sup>18</sup> Yeung SSY, Trappenburg MC, Meskers CGM, Maier AB, Reijnierse EM. Inadequate energy and protein intake in geriatric outpatients with mobility problems. Nutr Res. 2020. doi:10.1016/j.nutres.2020.09.007
- <sup>19</sup> Geirsdottir OG, Arnarson A, Ramel A, Jonsson P V, Thorsdottir I. Dietary protein intake is associated with lean body mass in community-dwelling older adults. Nutr Res. 2013;33(8):608-612. doi:10.1016/j.nutres.2013.05.014
- <sup>20</sup> McLean RR, Mangano KM, Hannan MT, Kiel DP, Sahni S. Dietary Protein Intake Is Protective Against Loss of Grip Strength Among Older Adults in the Framingham Offspring Cohort. Journals Gerontol - Ser A Biol Sci Med Sci. 2016;71(3):356-361. doi:10.1093/gerona/glv184
- <sup>21</sup> Isanejad M, Mursu J, Sirola J, et al. Dietary protein intake is associated with better physical function and muscle strength among elderly women. Br J Nutr. 2016;115(7):1281-1291. doi:10.1017/S000711451600012X



- <sup>22</sup> Cuesta-Triana F, Verdejo-Bravo C, Fernández-Pérez C, Martín-Sánchez FJ. Effect of Milk and Other Dairy Products on the Risk of Frailty, Sarcopenia, and Cognitive Performance Decline in the Elderly: A Systematic Review. Adv Nutr. 2019;10(suppl\_2):S105-S119. doi:10.1093/advances/nmy105
- <sup>23</sup> Lana A, Rodriguez-Artalejo F, Lopez-Garcia E. Dairy consumption and risk of frailty in older adults: A prospective cohort study. J Am Geriatr Soc. 2015;63(9):1852-1860. doi:10.1111/jgs.13626
- <sup>24</sup> Otsuka R, Tange C, Tomida M, et al. Dietary Factors Associated with the Development of Physical Frailty in Community-Dwelling Older Adults. J Nutr Heal Aging. 2019;23(1):89-95. doi:10.1007/s12603-018-1124-3
- <sup>25</sup> Learn What Osteoporosis Is and What It's Caused by. https://www.nof.org/patients/what-is-osteoporosis/. Accessed December 10, 2019.
- <sup>26</sup> US Office of the Surgeon General. Bone Health and Osteoporosis. Office of the Surgeon General (US); 2004. http://www.ncbi.nlm.nih.gov/ pubmed/20945569. Accessed April 4, 2018.
- <sup>27</sup> Shi Y, Zhan Y, Chen Y, Jiang Y. Effects of dairy products on bone mineral density in healthy postmenopausal women: a systematic review and metaanalysis of randomized controlled trials. Arch Osteoporos. 2020;15(1):1-8. doi:10.1007/s11657-020-0694-y
- <sup>28</sup> Hidayat K, Du X, Shi BM, Qin LQ. Systematic review and meta-analysis of the association between dairy consumption and the risk of hip fracture: critical interpretation of the currently available evidence. Osteoporos Int. 2020;31(8):1411-1425. doi:10.1007/s00198-020-05383-3
- <sup>29</sup> Ong AM, Kang K, Weiler HA, Morin SN. Fermented Milk Products and Bone Health in Postmenopausal Women: A Systematic Review of Randomized Controlled Trials, Prospective Cohorts, and Case-Control Studies. Adv Nutr. 2020;11(2):251-265. doi:10.1093/advances/nmz108
- <sup>30</sup> Fabiani R, Naldini G, Chiavarini M. Dietary patterns in relation to low bone mineral density and fracture risk: A systematic review and meta-analysis. Adv Nutr. 2019;10(2):219-236. doi:10.1093/advances/nmy073
- <sup>31</sup> Malmir H, Larijani B, Esmaillzadeh A. Consumption of milk and dairy products and risk of osteoporosis and hip fracture: a systematic review and Metaanalysis. Crit Rev Food Sci Nutr. 2020;60(10):1722-1737. doi:10.1080/10408398.2019.1590800
- <sup>32</sup> Wallace TC, Bailey RL, Lappe J, et al. Dairy intake and bone health across the lifespan: a systematic review and expert narrative. Crit Rev Food Sci Nutr. 2020. doi:10.1080/10408398.2020.1810624
- <sup>33</sup> Feskanich D, Meyer HE, Fung TT, Bischoff-Ferrari HA, Willett WC. Milk and other dairy foods and risk of hip fracture in men and women. Osteoporos Int. 2018;29(2):385-396. doi:10.1007/s00198-017-4285-8
- <sup>34</sup> Aslam H, Holloway-Kew KL, Mohebbi M, Jacka FN, Pasco JA. Association between dairy intake and fracture in an Australian-based cohort of women: A prospective study. BMJ Open. 2019;9(11). doi:10.1136/bmjopen-2019-031594
- <sup>35</sup> Wallace TC, Jun S, Zou P, et al. Dairy intake is not associated with improvements in bone mineral density or risk of fractures across the menopause transition: data from the Study of Women's Health Across the Nation. Menopause. 2020;27(8):879-886. doi:10.1097/GME.00000000001555
- <sup>36</sup> Machado-Fragua MD, Struijk EA, Caballero FF, et al. Dairy consumption and risk of falls in 2 European cohorts of older adults. Clin Nutr. 2020;39(10):3140-3146. doi:10.1016/j.clnu.2020.01.025
- <sup>37</sup> Matía-Martín P, Torrego-Ellacuría M, Larrad-Sainz A, Fernández-Pérez C, Cuesta-Triana F, Rubio-Herrera MÁ. Effects of Milk and Dairy Products on the Prevention of Osteoporosis and Osteoporotic Fractures in Europeans and Non-Hispanic Whites from North America: A Systematic Review and Updated Meta-Analysis. Adv Nutr. 2019;10(suppl\_2):S120-S143. doi:10.1093/advances/nmy097